International Academic Institute for Science and Technology

International Academic Journal of Accounting and Financial Management
Vol. 5, No. 1, 2018, pp. 28-34.

ISSN 2454-2350

International Academic Journal of Accounting and Financial Management

www.iaiest.com

Studies on Symptomological and Economic Parameters of Silk Cocoons of *Bombyx Mori* Inoculated with *Beauveria Bassiana* (Bals.) Vuill.

Seema K. Dongare[a], Priti M. Gaikwad[b], Shubhangi S[c]. Pawar[d], Vitthalrao B. Khyade[e]*

Department of Zoology, Shardabai Pawar Mahila Mahavidyalaya, Shardanagar; Tal. Baramati; Dist. Pune – 413115 (India).
**: Author for correspondence*

Abstract

The muscardine disease in the larval instars of silkworm is caused by fungus, *Beauveria bassiana* (Bals) Vuill. The Beauveria *bassiana* (Bals) Vuill is one of the most destructive pathogen of silkworm *Bombyx mori* L and it is common in all sericulture zones of the world. The fungal disease is common during winter and rainy seasons. White muscardine was the first disease described in insects caused by a microorganism. This fungus was named eventually *Beauveria bassiana* in honor of Agostino Bassi, sometimes called de Lodi (25 September 1773 – 8 February 1856), Italian entomologist, botanist and bacteriologist recognized as the "Father of Insect Pathology". *Beauveria bassiana* infection influenced the growth and development of silkworm larvae and ultimately the economical cocoon characters like matured larval weight, cocoon weight, shell weight, shell percentage, filament length, non-breakable filament length, number of breaks and denier. Significant reduction of matured larval weight (2.08 g), cocoon weight (0.72 g), shell weight (0.09 g), shell ratio (12.80%), filament length (478.9 m), non-breakable filament length (120.8 m) and more number of breaks (4.1) and higher denier (2.62 d) was recorded in the experimental silkworm compared to control.

Keywords: Beauveria Bassiana, Bombyx Mori, Symptomological Changes Cocoon Yield Parameters

Received:18 January 2018/Accepted:13 April 2018/Published:27 June 2018

Introduction:

Obviously, the success of sericulture industry primarily depends on the successful harvest of cocoon crops. The major problem of sericulture in a tropical country like India is the high incidence of diseases. The major diseases affecting mulberry silkworm are muscardine (fungal disease), flacherie (bacterial diseases) grasserie (viral diseases) and pebrine (protozoan disease). Silkworm crop loss occurs in all the silkworm growing areas of the world, but the type of severity varies. They differ from region to region, crop to crop and even from farmer to farmer.

Leaving aside minor variations, it has been found that crop loss is generally more in the tropics than in the temperate regions. The magnitude of the disease damage is on the higher side in India. It is a general observation that out of 5-6 crops per year, two are usually lost due to diseases and other reasons and even the successful once are partially lost. Thus the frequent outbreak of diseases is one of the main handicaps for the progress of sericulture industry. The reasons attributed, are poor hygienic conditions and continuous silkworm rearing all-round the year, the climatic conditions that favour faster multiplication of disease causing germs. The accumulated germs under favourable conditions become active and cause the outbreak of diseases. *Beauveria bassiana* (Bals) Vuill is one of the most destructive fungal pathogen of silkworm *Bombyx mori* L causing white muscardine disease, which is common in all sericulture zones of the world. The fungal disease is common during winter and rainy seasons. White muscardine was the first disease described in insects caused by a microorganism. Bassi (1835) demonstrated the contagious nature of the disease and identified the cause as a fungus and suggested measures to control the disease (Ainsworth, 1956). The fungus was named after Agostino Bassi Vuillemin in his honour as *Betrytis bassiana* by Balsamo and created a genus *Beauveria bassiana* (Balsamo) Vuillemin. It was a first microorganism to be recognized as a disease causing agent (Bassi, 1835). This disease wiped out the entire sericulture industry in Italy and France during 1920 1925. In India 10 40 percent of loss has been accounted for white muscardine in total loss due to diseases (Janakiraman, 1961; Ayuzawa, 1972; Chandrasekharan and Nataraju, 2008). In the year 2005, heavy rain in major silk producing states of Karnataka, Tamil Nadu and Andhra Pradesh has led to a widespread outbreak of muscardine among the silkworm cocoon crops, threatening the silk output in the country. In this backdrop, the study was carried out to examine the symptological changes in *Bombyx mori* during the progress of the disease and its influence on economical characters of the cocoons.

Materials and Methods

PM x CSR2 was selected for the study and the silkworms were reared under optimum conditions with meticulous coordination of many activities such as maintenance of mulberry garden, preparation and disinfection of the rearing room and appliances, procurement and handling of silkworm eggs and incubation, young and late age silkworm rearing, general hygiene, moulting care, mounting, spinning, harvesting of cocoons *etc.* (Dandin *et al.*, 2003). On the 1st day of the fifth instar, the larvae were inoculated by dipping in sublethal concentration of fungal conidia spore suspension (2.15 x 106 conidia spores/ ml @ 50 ml/100 worms for 45 Sec) and larvae treated with double distilled water were used as control. From the day of inoculation to the end of the 5th instar the silkworms were kept under continuous surveillance to examine the symptomplogical changes during the development of fungal pathogen *Beauveria bassiana* and taken photographs. By the end of the fifth instar the silkworm showed the symptoms of spinning i.e., cessation of feed, body shrinkage, translucent light yellow colour skin etc.

Received:18 January 2018/Accepted:13 April 2018/Published:27 June 2018

These larvae are known as matured worms, the weight of the worms was recorded. These worms were transferred to mountages to spin the cocoons. After completion of the spinning process, the cocoons were harvested on the 6th day of mounting and then these cocoons were taken for the assessment of various economic parameters viz., cocoon weight, shell weight, shell ratio, filament length, non-breakable filament length, number of breaks and denier. The qualitative and quantitative parameters of cocoons were determined by following the methods as given by Sonwalkar (1993).

Matured larval weight

This character indicates the healthy and robust disposition of the larvae. It is a mean of the weights of 10 randomly selected fully matured larvae.

Cocoon weight

Cocoon weight indicates the approximate quantity of raw silk that could be reeled from the cocoons (Mahadevappa *et al.,* 2000). In the present experiment a sample is drawn from each replication comprising around 10 cocoons. The sample drawn represents the entire quality of each replication. Individual cocoon weight was taken from each sample of 10 cocoons and
Mean cocoon weight was calculated. The weights were taken in gram units.

Shell weight

This economic trait represents the total quantity of silk in a cocoon. Average single shell weight was calculated from 10 shells used for the assessment of cocoon weight.

Shell ratio

It denotes the total amount of silk available in a single cocoon and is expressed in percentage. It is calculated by dividing the shell weight reading by cocoon weight reading. Quotient thus obtained was multiplied by hundred to obtain the shell ratio or shell percentage in entire cocoon.

Filament length

It is the total length of silk filament, unwound from a single cocoon measured in meters. In the present experiment a sample is drawn from each replication comprising around 10 cocoons. Ten cocoons were cooked and reeled on an eprouvette with a circumference of 1.125m and the mean value of filament length in meters was calculated as per the standard formula (Sonwalkar, 1993).
Filament length (m) = Number of rotations x Circumference of the wheel (1.125 m) Non-breakable

Filament length and Number of Breaks

It is the average length of filament that can be unwound from the cocoons without a break. Non-breakable filament length was calculated by using the formula as given by Sonwalkar (1993) and the number of breaks were recorded.
[Average Non-breakable filament Length = Total filament length ÷ (1 + Number of Breaks)].

Received:18 January 2018/Accepted:13 April 2018/Published:27 June 2018

Denier

Denier scale is the weight in grams of 9,000 meters of the filament. Denier represents the size of the yarn. It was calculated using the formula as given by Sonwalkar (1993).

Denier Scale = [Weightof filament(g) ÷ Lengthof filament(m)] x 9000

Variation in the denier of the cocoon silk filament will ultimately determine the uniformity and quality of raw silk yarn reeled.

Statistical analysis

All the experimental data recorded from the 4 replicates have been subjected to statistical analysis by following T-test.

Results and Discussion

Symptomological Changes

No conspicuous symptoms were noticed immediately after inoculation of fungal pathogen *Beauveria bassiana*. But gradually an infected silkworm became inactive, sluggish, stopped the feed and remained underneath the mulberry leaves. After 48 hours, the infected worms started to vomit digestive juice and later, the worms gradually became stiff and the movement of the worms was very much restricted and colour of the body changed to brown colour with oily specks. Initially the oily specks are in small in size with the advancement of the age, the size and number of the oil specks was enhanced (Photo. 3 & 5). Then the silkworm body became soft, pliable and later stiff and hard. Nearly on 7th or 8th day of the infection white efflorescence noticed near intersegment region, spiracles, then complete body was covered with the white mycelia and finally conidia developed on the body (Photo.4). The mummified cadaver became brittle and breaks into pieces when dropped from a certain height. Infected worms fail to spin the cocoons, but those which spin form flimsy cocoons (Photo.6). Cocoons formed by these infected worms were smaller and lighter in weight and the worms not emerged as moths.

Economical cocoon parameters

The overt changes observed in the economical traits of cocoon in *Beauveria bassiana* infected silkworm with reference to control are shown in Table-1 and Graphs-1-8. The results of the present study clearly indicated that *Beauveria bassiana* infestation influences the growth and
development of silkworm and the economical cocoon parameters. Significant reduction of matured larval weight (2.08 g), cocoon weight (0.72 g), shell weight (0.09 g), shell ratio (12.8%), filament length (478.9 m), non-breakable filament length (120.8 m) and more number of breaks (4.1) and higher denier (2.62 d) was recorded in the experimental silkworm larvae. In contrast, to the experimental larvae significantly higher values were recorded in the majority of the economical characters such as matured larval weight (2.58 g), cocoon weight (1.44 g), shell weight (0.24 g), shell ratio (16.43%), filament length (671.6 m), non-breakable filament length (640.6 m) and less number of breaks (0.2) and lower denier (2.2 d) in control.

Received:18 January 2018/Accepted:13 April 2018/Published:27 June 2018

Appropriate matured larval weight is an indicator to measure the health of silkworm and in turn to obtain good quality of cocoon. Cocoon weight is an important commercial character used to determine the amount of raw silk that can be obtained. The cocoon price is determined based on the weight of the cocoon. The shell weight is more important than the cocoon weight since it is
The shell that yields the silk for reeling. Thus, higher the weight of the shell, greater will be the silk yield. Shell percentage fairly indicates the quantity of raw silk that can be reeled from the cocoons and also helps in estimating renditta and thereby fixing a proper price for the cocoons. The silk filament length indicates the reelable length of silk filament from a cocoon. Reeling
Performance is better with cocoons having a larger filament length, which reduces the number of feeding ends. This results in increase of production, quality and also raw silk percentage of the cocoon. Generally, a longer non-breakable filament with less number of breaks, higher is the reelability. This indicates that higher the reelablity percentage higher is the raw silk quality.
The fineness of cocoon filament is expressed by size i.e. denier (Sonwalkar, 1993; Mahadevappa *et al.*, 2000). Reduction in the matured larval weight may be due to the consequence of fungal
Infection that leads to the decrease in food consumption, digestion, relative consumption rate, efficiency of conversion of ingested food in fifth instar of *Bombyx mori* infected with *Beauveria bassiana*. Cai (1989) analyzed the fungus growth and proliferation in silkworm body after 12 to 24 h post- inoculation of the *B. bassiana* conidium. The healthy growth and development of the silkworm is directly related to economical cocoon traits. It is well supported by Lakshmi Velide *et al.* (2013). She noticed poor cocoon characters from pebrine infected tropical tasar silkworm
Antheraea mylitta.

The reduction in economical cocoon characters could be attributed to loss of appetite, lethargic conditions and physiological stress induced by the fungal pathogen *Beauveria bassiana* in 5th instar silkworm larvae. Jhansi Lakshmi (2003) observed the reduction of economical parameters due to *Beauveria bassiana* infection and attributed that it may be due to the decline in the synthesis of silk proteins and the direct or indirect effect of *Beauveria bassiana* on the growth and development of silk gland of the silkworm *Bombyx mori*. More number of breakages and higher denier were noticed in experimental cocoons with reference to healthy ones. It may be
Assumed that the physiological and biochemical stress induced by a fungal pathogen caused to exude uneven amounts of silk fluid in lumps. Rath *et al* (2003) have reported the decrease in shell weight in *Antheraea mylitta* larvae infected with *Nosema* sp. Rath and Sinha (2005) studied
On the parasitization of fifth instar larvae of *Antheraea mylitta* by Uzifly have reported the decrease in cocoon weight and shell weight in the infected larvae.

Silkworm *Bombyx mori* is completely domesticated economic insect and *Beauveria bassiana* is an aggressive parasite. Therefore, the disease in silkworm results in a drastic reduction in the growth and development and qualitative and quantitative cocoon yield parameters. This ultimately has a direct effect on the sericulture community of the country due to reduced returns, in turn affecting the economy of the country, since silk as the end product of sericulture industry is a good source of foreign exchange for the country. Therefore, it is the prime responsibility of

Received:18 January 2018/Accepted:13 April 2018/Published:27 June 2018

Sericulture scientists and sericulture farmers to explore many more strategies to prevent/control the occurrence of the disease to enhance the qualitative and quantitative parameters of cocoon crops.

Acknowledgement

Authors are highly grateful to Agriculture Development Trust, Shardanagar (Baramati) for valuable help, providing facilities at the sericulture unit of Malegaon sheti farm & laboratory facilities at Shardabai Pawar Mahila College to carry out the experimentations.

References

Ainsworth, G.C. 1956. Agostino Bassi, 1773-1856. Nature, 177: 255 257.

Ayuzawa, C. 1972. Studies on the infectious flacherie of the silkworm, Bombyx mori L. Purification of the virus and some of its properties. J. Seric. Sci. Tokyo., 41: 338.

Bassi, A. 1835. Del mal Del Segno calcinaccio moscardino malatta che affiggei bachidaseta parte 1. Teorica Tip. Orcesi, Lodi.

Cai, J. 1989. By scanning electron microscope, White cadaver skin bacteria invade the body of silkworm [J]. Chin. J. Nature, 12(11): 857 858.

Chandrasekharan, Nataraju, B. 2008. Studies on white muscardine disease of mulberry silkworm, Bombyx mori L. in India. Indian J. Seric., 47(2): 136 154.

Dandin, S.B., Jayant Jayaswal, Giridhar, K. 2003. Hand Book of Sericulture Technologies. Central Silk Board,Bangalore.

Janakiraman, A.T. 1961. Diseases affecting the Indian silkworm races. J. Silkworm, 13: 91 101.

Jhansi Lakshmi, V.V.N.S. 2003. Ultrastructural studies on tissues of the silkworm, Bombyx mori L infected with Beauveria bassiana (Balsamo) Vuillemin. Thesis submitted to Sri Padmavati Mahila Universiy, Tirupati. Pp. 58 70.

Lakshmi Velide, Bhagavanulu, M.V.K., Purushotham Rao, A. 2013. Study on impact of parasite (Nosema species) on characters of tropical tasar silkworm, Antheraea mylitta drury J. Environ. Biol., 34: 75 78.

Mahadevappa, D., Halliyal, V.G., Shankar, D.G., Ravindra Bhandiwad. 2000. Mulberry silk reeling technology. Oxford & IBH Publishing, New Delhi.

Rath, S.S., Prasad, B.C., Sinha, B.R. 2003. Food utilization efficiency in fifth instar larvae of Antheraea mylitta (Lepidoptera: Saturniidae) infected with Nosema sp. and its effect on reproductive potential and silk production. J. Invert. Pathol., 83: 1 9.

Rath, S.S., Sinha, B.R.R.P. 2005. Parasitization of fifth instar tasar silkworm, Antheraea mylitta by the uzifly Blepharipa zebina; a hostparasitoid interaction and its effect on hosts nutritional parameters and parasitoid development. J. Invertebr. Pathol., 88: 70 78.

Sonwalkar, T.N. 1993. Hand book of silk technology. Wiley Eastern ltd, New Delhi.

Received:18 January 2018/Accepted:13 April 2018/Published:27 June 2018

Table.1: Changes in economic cocoon parameters of silkworm *Bombyx mori* L. inoculated with fungal pathogen *Beauveria bassiana* (Bals.) Vuill. With reference to control.

Paramateres Groups ↓	**Matured larval Weight (g)** →	**Cocoon weight (g)**	**Shell Weight (g)**	**Shell Ratio (%)**	**Filament Length (m)**	**Non-Breakable Filament Length (m)**	**Number of Breaks**	**Denier (d)**
Control	2.58 (±0.07)	1.44 (±0.06)	0.24 (±0.03)	16.43 (±0.12)	671.6 (±18.33)	640.6 (± 14.08)	0.2 (± 0.42)	2.20 (±0.12)
Diseased (Inculated)	2.08 (±0.07) ****	0.72 (±0.17) ****	0.09 (±0.01) ****	12.80 (±0.10) ****	478.9 (±15.74) ****	120.8 (± 22.42) ****	4.1 (± 2.42) ****	2.62 (±0.07) ****

Mean±Standard Deviation; ****P 0.001

Received:18 January 2018/Accepted:13 April 2018/Published:27 June 2018