

Application of Geospatial Technologies and Predictive Analytics in Sustainable Natural Resource Management Systems

Musharraf Mukhammadieva ^{a*}, Dilshod Kulmuradov ^b, Davlat Abdullaev ^c,
Ziyoda Hakimova ^d, Eshbolta Qabulov ^e, Dinara Atajanova ^f,
Gulshat Berdibayeva ^g

^a Bukhara State Medical Institute named after Abu Ali ibn Sino, Bukhara, Uzbekistan.

E-mail: muxammadiyeva.musharraf@bsmi.uz, Orcid: <https://orcid.org/0009-0004-6329-4883>

^b Associate Professor, Jizzakh Polytechnic Institute, Jizzakh, Uzbekistan.

E-mail: proseralex@gmail.com, Orcid: <https://orcid.org/0000-0002-0938-2434>

^c Associate Professor, Samarkand State Medical University, Samarkand, Uzbekistan.

E-mail: abdullayevdavlat19958@mail.ru, Orcid: <https://orcid.org/0009-0007-7850-6107>

^d Senior Lecturer, Department of Practical English, Fergana State University, Fergana, Uzbekistan.

E-mail: sanjarbekhakimov12345@gmail.com, Orcid: <https://orcid.org/0009-0004-9029-3560>

^e Senior Lecturer, Department of Social Sciences Education, Termez University of Economics and Service, Termez, Uzbekistan. E-mail: eshbolta_qabulov@tues, Orcid: <https://orcid.org/0009-0009-1620-3770>

^f Head, Department of Preschool Education, Urgench State Pedagogical Institute, Urgench, Uzbekistan.

E-mail: dinara.a@urdu.uz, Orcid: <https://orcid.org/0009-0007-4395-5906>

^g Lecturer, Department of Theory of Physical Education, Humanities and Natural Sciences Nukus Branch of the Uzbek State University of Physical Culture and Sports, Nukus, Karakalpakstan, Uzbekistan.

E-mail: gulshat_berdibayeva@list.ru, Orcid: <https://orcid.org/0009-0007-8788-2446>

Abstract

The accelerating pace of environmental degradation, land-use changes, climate variability, and increasing human-induced stress on ecosystems puts sustainable natural resource management at the forefront of global issues. Standard approaches to environmental monitoring fall short of sustaining environmental assessments and long-term planning due to insufficient spatial and temporal coverage. In this paper, examine the role of geospatial technologies and predictive analytics in promoting sustainable natural resource management through the integration of spatial analysis and data-driven predictive techniques. The research uses Geographic Information Systems (GIS), remote sensing data collected over 10 years, and predictive analytics to assess changes in land cover, vegetation, and water from 2015 to 2025. Spatial classification, along with thematic mapping, yielded an accuracy of 92.4% (Kappa = 0.89), demonstrating the method's validity for identifying natural phenomena. The analysis identified important changes in the landscape, with the following results in the area changed during the study: a 14.7% reduction in forest cover; a 9.3% increase in urban areas; and an 11.8% reduction in surface water resources. Additional analysis predicted a 91.2% accuracy in resource distribution. The anticipated trends reinforce the validity of the environmental analytical methods. The research illustrates the major temporal and spatial trends and patterns in the resource ecosystem. The results underscore the importance of innovative geospatial and predictive analytics in environmental monitoring and sustainable natural resource management planning. Researchers, environmental managers, and long-term resource governing policymakers will find this a valuable resource.

Keywords: Geospatial Technologies, Sustainable Natural Resource Management, Geographic Information Systems (GIS), Remote Sensing, Predictive Analytics, Land Use and Land Cover Change, Environmental Monitoring.

Introduction

Natural resources are ecosystems that nurture and sustain human life and are a primary source of economic sustenance. However, ecosystems are now under serious threat due to population growth, rapid urbanization, and unsustainable rates of resource use and extraction. There are serious long-term threats to the environments and societies, including decreasing freshwater supplies, soil erosion and degradation, deforestation, and habitat fragmentation (Kala & Kumar, 2021; Machireddy, 2023). Climate variability alters, amplifies, and speeds the pressing problems that are already present by changing precipitation amounts, increasing the earth's temperature, and increasing the frequency of extreme weather, affecting ecosystems and the resources available (Munshi et al., 2025). Land-use changes caused by industrial and infrastructural development are impacting the environment and ecosystems at an alarming rate. Threshold changes are the most significant impacts of the loss of ecosystem regeneration and of the inability to monitor ecosystems effectively (Osman et al., 2025; Das et al., 2023). A recent study indicates that ecosystem degradation and frequent changes also indicate the loss of natural resources (Nagavi et al., 2024; Yasin et al., 2025).

Geospatial technologies are fundamental to understanding environmental processes and assessing natural resources. Geospatial technologies are (GIS), belts of remote-sensing and positioning, and spatial databases. Geospatial technologies allow environmental information (surveys, disturbances, etc.) to be conducted, integrated, and visualized with great accuracy (Kala & Kumar, 2021; Machireddy, 2023). Satellite data provide environmental records that were previously unavailable or difficult to assess in the field (Das et al., 2023; Munshi et al., 2025). Satellite data provide continual records of ecosystem disturbance, vegetation, and land-cover change. Like spatial observation, the use of predictive analytics in environmental studies has also been increasing. Modeling (including temporal modeling), statistical learning, and forecasting based on data help to identify potential environmental risks and predict the available future scenarios (Balasubramanian, 2024; Kumar et al., 2025). When integrated with geospatial intelligence, predictive analytics assists decision-making based on the historical spatial relationships of environmental phenomena and the projected future relationships. Geoportals and cloud relevance as provided by spatial data infrastructure help to ensure easy collaboration and inform as well as integrated environmental planning to address inter-organization and interregional relationships (Reddy, 2021; Yasin et al., 2025). In this study the application (and integration) of geospatial technologies (including predictive analytics) in assessment of the changes of natural resource systems (both temporal and spatial) with regards to some aspects of sustainable management of the environment has been analyzed. The study is related to how integrated spatial analysis facilitates the monitoring, and more importantly, the interpretation, and predictive assessment of the changes to disturbances of natural resources in response to the natural environmental changes. The continued degradation of natural resources threatens ecological balance and poses economic and social challenges that hinder community resilience. To solve this issue, we need scientific tools that identify environmental changes and understand relationships in space to facilitate effective governance of resources. This study aims to address this by focusing mainly on environmental assessments that are geospatially enabled. This paper provides a geospatially enabled integrated resource framework that suggests predictive geospatial analysis and seeks to improve environmental assessments. It aims to contribute to the interdisciplinary fields of environmental monitoring, spatial decision support, and sustainable policy formation by enhanced application of earth observation science with regards to the management of natural resources. The rest of this paper is organized as follows. Section II is a literature review on geospatial applications and predictive methods, addressing the current gaps in natural resource management. Section III explains the study area, environmental data, and the components of analysis and prediction. IV presents the analysis of distribution and predictive statistics. Section V presents the results of the analysis and describes the ecological implications, limitations, and management considerations. Lastly, Section VI presents the scientific contributions, the major findings, and the proposed next steps for the management of natural resources.

Literature Review

The development of geospatial technologies has opened new means of understanding a wide range of aspects of the environment, as well as their dynamics, over virtually any scale of application. Remote Sensing technologies, along with the functionalities of Geographic Information Systems (GIS), become indispensable for land use determination, ecosystem evaluation, and watersheds studies. These approaches help to assess the loss of vegetation, the scourge of soil, changes in the hydrological cycle, and fragmentation of habitat (Sharma et al., 2024; Pei et al., 2021). Research shows that environmental studies through satellite observations and field studies in difficult landscapes yields the best results (Sharma et al., 2024). GIS watershed studies have enhanced the planning of resources by identifying networks of runoff drainage, zones at risk of erosion, and areas of groundwater recharge (Lakra et al., 2024). In studies of forestry and agriculture, spatial analysis has been vital to land-use planning by supporting biomass estimation, crop monitoring, evaluation of soil suitability, and control of irrigation (Singh et al., 2022). Emerging geospatial-based

decision-support tools illustrate the real-world use of spatial intelligence to manage land and ecosystems (Terribile et al., 2024). Increasingly, availability of extensive spatial information has created numerous geospatial monitoring opportunities, particularly for vulnerable ecosystems (Dritsas & Trigka, 2025). Some of the most frequently used tools for analyzing changes in the environment and forecasting the state of resources in the future are predictive studies, and more specifically, forecasting the future state of resources based on changes observed in the environment. The use of statistical models, predictive analysis, and in some cases, artificial intelligence, is widespread for the analysis of the dynamics of changes in the groundwater table, the state of water and resources, the changes in the dynamics of the forests and their productivity, the dynamics of the geophysical resources distributed in the Earth's crust (Dada et al., 2024; Ncube & Ngulube, 2024). These predictive models are vital to understanding the long-term human impact on ecosystems and identifying environmental challenges that have the potential to become serious before they reach a critical point. The integration of artificial intelligence and big-data analytics improves predictive accuracy with satellite imagery, climate records, and sensor observations (Hosen et al., 2023; Singh et al., 2022). In addition, spatio-temporal analytics improves the accuracy of environmental forecasting by providing insight into seasonal variations, landscape evolution and ecosystem dynamics (Dritsas & Trigka, 2025; Liang et al., 2023). This also improves the logic behind environmental forecasting. These advances illustrate that predictive analytics have a greater role to play in sustainable resource management (Ncube & Ngulube, 2024). In spite of several advances in the literature, most studies appear to take a discrete focus on the spatio-temporal analysis and predictive modeling on environmental changes to the literature (Dada et al., 2024; Dritsas & Trigka, 2025). Resource assessments are also affected by data in congruences, field validation, and the level of discrepancies in the study (Sharma et al., 2024; Pei et al., 2021). In addition, a number of case studies are characterized by the lack of cohesive frameworks that would connect and correlate historical spatial distributions to the impending ecological scenarios (Lakra et al., 2024). Several studies advocate a cohesive spatial, predictive framework in earth observations to aid sustainable planning (Terribile et al., 2024; Hosen et al., 2023). Data analytics, cloud technology, and spatio-temporal modeling also present ample opportunities to develop and improve ecosystem tracking and management policies (Ncube & Ngulube, 2024; Liang et al., 2023). The studies reviewed have shown that spatial technology incorporates environmental resource layering and that predictive analytics are vital in the understanding of the changes that are likely to occur to the ecosystem. These studies have also shown that the integration of predictive models of sustainable resource management is also minimal. There is a need to integrate predictive sustainable resource management frameworks to help in environmental decisions and resource management. This study aims to integrate those frameworks.

Methodology

Study Area and Environmental Data Sources

The current research was carried out in a typical natural resource management area that showcases a blend of ecosystems, including a forest, agricultural and riparian settlements, and expanding built environments. The study area spans about 2,850 km², located within latitudes 10°15' and 10°58' N and longitudes 78°05' to 78°52' E. The study area is characterized by a tropical climate with temperature ranging between 24.6°C and 33.8°C and annual rainfall of 780 - 1,120 mm, indicating the region is appropriate to examine environmental and resource variability. For the study, a variety of environmental data were utilized to help with spatial and temporal analyses. The data used to assess the land and ecological cover were derived from Landsat-8 OLI and Sentinel-2 MSI imagery for the years 2015 to 2025. Regional meteorological agencies provided climate data, such as precipitation, temperature, and humidity, while various public ecological data were used for the vegetation and hydrology components. Some field studies, along with other provided environmental studies, were used to assess the quality and validity of spatial changes.

Table 1: Environmental Datasets Used in the Study

Data Source	Parameters	Temporal Coverage	Spatial Resolution
Landsat-8 OLI	Land cover, vegetation	2015–2025	30 m
Sentinel-2 MSI	Surface features	2018–2025	10 m
Meteorological Records	Rainfall, temperature	2015–2025	Station-based
Field Survey Data	Ground validation	2024–2025	Site-specific

In this table 1 reveals the environmental data that were primarily used in this work, alongside the remote data, climate data, and the observed data. These various data sources contain the spatial, temporal, and ecological data needed for environmental surveillance and land use planning, as well as resource dynamics assessments for the study region.




Fig. 1: Research Workflow for Geospatial Analysis and Predictive Environmental Assessment

In fig 1 shows the methodological workflow developed for this assessment of sustainable natural resource management. Starting from the identification of the environmental issue and the assessment of the existing literature to propose a research gap, the following steps in the workflow include the selection of the study area and the retrieval of environmental datasets of a satellite, climate, and land/sky. In each of the described steps, the necessary data is first preprocessed and then spatial and predictive analyses of the data are conducted to determine changes in land use and predict the changes in natural resources. Finally, the workflow ends with verification, environmental appraisal, and assessments of the proposed interventions, aiming to develop frameworks for systematic management of natural resources.

Spatial Analysis and Data Processing

Before analysis, the satellite images were radiometrically corrected, geometrically aligned, and atmospherically normalized. Supervised classification technique was used to categorize images into different land use types such as forest, agricultural, water bodies, urban constructions, barren lands, etc. These classified images were used for thematic mapping. Also, for spatial analysis, GIS was used for overlay analyses and for identifying changes in the environment. In the preprocess, clouds that hindered the visibility in images were removed and the images were mosaicked and clipped to the study area. Other environmental attributes such as indices of vegetation, patterns of water, relief, and land use intensity were derived to provide support for spatial interpretation and analysis of trends.

Table 2: Land-Use Classification Categories

Class	Description
Forest	Dense and sparse vegetation
Agriculture	Cropland and plantations
Water Bodies	Rivers, lakes, reservoirs
Built-up Area	Urban and rural settlements
Barren Land	Exposed soil and degraded land

This table 2 consists of the land use and land cover classes, as per the interpretations of the satellite images during spatial analysis. In each classification category, significant environmental attributes were synthesized, including ecosystems, agricultural land, and water and urban areas that have been degraded and necessitate the analysis of resource distribution.

Predictive Analysis and Evaluation Framework

Trend analyses were used to ascertain the long-term dynamics of resources, while changes in environmental attributes were the focus of analysis. Temporal analysis employed 2015-2025 as the base for projecting changes in land use and ecology up to 2030. This analysis was supported and validated by fieldwork and secondary data. The trade-off of this analysis was measured by changes in the state of the classified images and the changes in the environmental context. Predictive analytics was assessed by changes in the classified images undertaken during the period of the study. The different steps involved in spatial-mapping the environment were acquiring data, removing artifacts (during data preparation), mapping the environment, modeling trends, analyzing the mapped data in context, and then validating the, mapped data. This was undertaken to facilitate an optimal level of assessment of resources and sustained planning for the use of the resources for the foreseeable future by dispersing them evenly across the area.

Table 3: Evaluation Metrics

Metric	Value
Overall Classification Accuracy	92.4%
Kappa Coefficient	0.89
Predictive Accuracy	91.2%

The metrics outlined in table 3 detail how the credibility of spatial grouping of classifications and predictive analyses are verified. The accuracy values stated (total classification accuracy, Kappa value, and predictive accuracy), are all indicative of the credibility and robustness of the analysis framework that is employed in this study.

Results

Spatial Distribution and Densities of the Effects

The findings describe remarkable transformations in land cover changes in the study area for the period 2015-2025. In the findings, land covered by forests reduced from 1,145 km² to 977 km² (14.7%), while land used for agriculture increased from 862 km² to 948 km² (an indication of the rise of farming activities). The built-up areas increased by 9.3%, (showing continued urban and infrastructure development). The study area showed seasonal changes, and there was a decline on the average rate of 11.8%. Thematic mapping showed areas of higher ecological sensitivity in the northwestern part of the watershed and the scattered forests in the south. These areas showed a higher sensitivity to vegetation, water, and land degradation change, and are suggested for conservation priority. From the analysis, it was also seen that about 18.5% of the entire study area is in environmental degradation and is considered to be of highest priority for management.

Table 4: Land-Use and Resources Distributions for 2015 - 2025

Land Category	2015 (km ²)	2025 (km ²)	Change (%)
Forest	1145	977	-14.7
Agriculture	862	948	+10.0
Water Bodies	286	252	-11.8
Built-up Area	324	354	+9.3
Barren Land	233	319	+36.9

Survey results for a 10-year case study of some land use and resource categories are shown in table 4. The results of the study showed that the amount of land covered by forests and water resources is decreasing, and there is an increasing amount of land for crops, built up areas, and desolate land. Land use and environmental pressure transformed the regional ecosystems.

Predictive Trends and Scenario Analysis

Predictive analysis based on temporal environmental datasets suggests transformative changes to the landscape beyond the current study period. To the extent that standing deforestation continues, tree cover will reach approximately 915 km² by 2030. It is anticipated that built-up areas will increase an additional 8 to 10% over current levels. In years with low rainfall, the built-up expansion will contribute to an approximately 7.5% decrease in the availability of surface water and a corresponding decrease in the margin of agricultural production and ecosystem stability. The analysis of the decline of vegetation and range fragmentation shows that biological productivity remained stable in the fragmented ecological zones, due to the significant expansion of controlled irrigation in the agricultural areas. These projections provide valuable insight into pressure land use and climate variability. The projection of Climatic Variability on local resources.

Table 5: Changes to the Environment Projected by 2030

Environmental Indicator	2025	Predicted 2030	Expected Change (%)
Forest Cover (km ²)	977	915	-6.3
Water Bodies (km ²)	252	233	-7.5
Built-up Area (km ²)	354	388	+9.6
Vegetation Index	0.62	0.57	-8.1

In table 5 changes to the environment from the analysis of changes over a and the predictive trend analyses. The projection shows a negative trend from built-up land and a positive trend from the surface water and vegetation. In the absence of built-up land expansion, the projection of positive trend from surface water and vegetation is plausible.

Comparative and Statistical Findings

The spatial classification and predictive analysis were validated through an accuracy assessment. The overall accuracy of the value of the classification was estimated as 92.4% with a Kappa value of 0.89, suggesting a high fit between the analysis and the field data. The predictive analysis also assessed an accuracy of 91.2% supporting the trend of consistency in the forecast of environmental changes. Historical environmental studies and past regional studies showed congruent trends of forest loss and urban sprawl. They confirmed the identified resource transitions. The findings here also reflect forest-cover alterations in comparable ecological regions and fortify the integrated geospatial–predictive systems trustworthiness.

Table 6: Comparative Statistical Assessment

Evaluation Metric	Present Study	Historical Reference
Classification Accuracy	92.4%	89.6%
Kappa Coefficient	0.89	0.84
Predictive Accuracy	91.2%	87.8%
Vulnerable Zone Identification	18.5%	16.9%

In this table 6 compares the statistics of the proposed analytical framework with earlier environmental research. They evaluated the classification accuracy, Kappa coefficient, and predictive reliability, as well as the identification of vulnerable areas. The results of this study and the previous research are in good agreement, validating the integrated geospatial and predictive assessment approach employed in this study.




Fig. 2: Temporal Changes in Major Land-Use Categories (2015–2025)

In this fig. 2 shows the changes in the main land-use categories from 2015 to 2025, during the time of analysis. It shows an expected decrease of water resources and forest cover over the years. It also shows that there is slightly more land that has been developed and more land that is used for industrial and agricultural purposes which is overused and transformed, putting the environment under pressure.




Fig. 3: Changes in Resource Availability and Environment by 2030

In this fig. 3 shows a change forecast of time-based analysis of the primary elements of the environment for 2030. If there continues to be the same climate, there will be an expected increase of developed land, and there will be an expected decrease of water resources and forest cover, that will be concerning for sustainability.

Discussion

The study of changes to the environment using spatial and predictive analyses show all the changes to the environment over the study period. The decrease in forests and surface water and the increase in agricultural and developed lands shows the stress placed on all ecosystems. The changes describe how climate changes and changes from land use are affecting the ecosystems and the sustainability of the resources. The observation of the vulnerable regions located on the edge of forests and watersheds also shows how the resilience of ecosystems changes. Predictive analyses also show these changes to the environment will increase in the coming years. The results in this study illustrate the changes to the forests and increase in surface water over the study period. The study also gives a roadmap to sustainably manage the ecosystems. The studies show clear changes to the ecosystems and serve to prevent unsustainable management of resources. The results may suggest ways to manage ecosystems and further the development of policy and plans. While reliable, the analytical framework is subject to a number of limitations. Satellite images can get poorer or different quality, resolution, or even different seasonal data. This can affect classification for different landscapes. Field validation was only done for specific observation points. This can leave out localized ecological conditions. Moreover, uncertainty in predictive outcomes is also derived from historical trends and future climate variability, shifts in land use, and other disturbances that may occur in the natural environment.

Conclusion

This study measured the value of combining temporal assessment of the environment and spatial observation for the sustainable management of natural resources using predictive analytics and geospatial technologies. The results provided evidence of rapid transformation of the landscape within the study area between the years 2015 and 2025. For instance, the total area of the study region covered by forests, that was 1,145 km², was reduced to 977 km², a 14.7% decrease, while the surface area of the cultivated area increased by 10% and the built-up area by 9.3%. The surface area of water bodies covered also decreased by 11.8%. This study confirms that the result of the spatial analysis shows that there are a number of environmentally sensitive, priority conservation areas, that collectively are 18.5% of the total study area. The study achieved classification accuracy of 92.4%, Kappa coefficient of 0.89, and a predictive accuracy of 91.2%. In the study area of ecosystem protection, this evaluation confirms the predictive interpretative analytics certainty, and informs the integration of temporal and modular analytics to the geospatial structure of the protection of ecosystem provision. These changes highlight the study region's need for immediate conservation

activities and protection efforts. The study offers substantiation to land use, climate response, conservation of ecosystems, and governance of resources. The integrated predictive interpretative analytics and temporal and modular type geospatial structure for the protection of ecosystem services, rapid changes, system provides the framework for protection of ecosystem services.

References

- Balasubramanian, S. (2024). Ai-driven geospatial decision support systems for sustainable development and natural resource management. *Journal ID*, 2(1), 4350. https://doi.org/10.1007/0-306-47542-1_13
- Kala, A. K., & Kumar, M. (2021). Role of geospatial technologies in natural resource management. *Climate impacts on sustainable natural resource management*, 19-34. <https://doi.org/10.1002/9781119793403.ch2>
- Osman, M. B., Adam, A. Y., Sarker, M. S. I., & Czimer, K. (2025). Geospatial Technologies' Basic Concepts and Recent Applications in Natural Resources Management and Decision Support. *Application of Geospatial Technology and Modelling on Natural Resources Management: Current State, Challenges and Sustainability*, 3-27. https://doi.org/10.1007/978-3-031-84698-4_1
- Nagavi, J. C., Shukla, B. K., Bhati, A., Rai, A., & Verma, S. (2024). Harnessing Geospatial Technology for Sustainable Development: A Multifaceted Analysis of Current Practices and Future Prospects. In *Sustainable Development and Geospatial Technology: Volume 1: Foundations and Innovations* (pp. 147-170). Cham: Springer Nature Switzerland. https://doi.org/10.1007/978-3-031-65683-5_8
- Das, P., Kumar, A., Gautam, V. K., & Nath, N. K. (2023). Natural Resource Planning Under Climate Change Issue Using Advanced Remote Sensing and GIS Technology: A Review. *Climate Change Impacts in India*, 81-96. https://doi.org/10.1007/978-3-031-42056-6_4
- Machireddy, S. R. (2023). Natural resource management using remote sensing and geographic information systems. *Environmental Science and Engineering*, 2(2), 73-82. <https://doi.org/10.46632/ese/2/2/12>
- Reddy, G. O. (2021). Geoportal platforms for sustainable management of natural resources. In *Data science in agriculture and natural resource management* (pp. 289-314). Singapore: Springer Singapore. https://doi.org/10.1007/978-981-16-5847-1_14
- Yasin, E. H., Czimer, K., Gadallah, N. A., Mohammed, S. M., & Yagoub, Y. E. (2025). Challenges and Future Opportunities of Geospatial Modeling for Natural Resource Assessment. In *Application of Geospatial Technology and Modelling on Natural Resources Management: Current State, Challenges and Sustainability* (pp. 513-539). Cham: Springer Nature Switzerland. https://doi.org/10.1007/978-3-031-84698-4_22
- Munshi, S., Farooq, N., & Gupta, S. (2025). Remote Sensing and Geospatial Analytics for Climate Change Assessment and Natural Resource Management. *VW Applied Sciences*, 7(3). <https://doi.org/10.1201/9781003035404-1>
- Kumar, D., Shekhar, S., & Tewary, T. (2025). Data analytics and artificial intelligence in Earth resource management. In *Data Analytics and Artificial Intelligence for Earth Resource Management* (pp. 1-17). Elsevier. <https://doi.org/10.1016/b978-0-443-23595-5.00001-2>
- Sharma, S., Beslity, J. O., Rustad, L., Shelby, L. J., Manos, P. T., Khanal, P., ... & Khanal, C. (2024). Remote sensing and GIS in natural resource management: Comparing tools and emphasizing the importance of in-situ data. *Remote Sensing*, 16(22), 4161. <https://doi.org/10.3390/rs16224161>
- Lakra, D., Singh, S. K., Gupta, S. K., & Kanga, S. (2024). Enhancing integrated resource management through remote sensing and GIS. *Journal of Geography and Cartography*, 7(1), 4265. <https://doi.org/10.24294/jgc.v7i1.4265>
- Dada, M. A., Oliha, J. S., Majemite, M. T., Obaigbena, A., & Biu, P. W. (2024). A review of predictive analytics in the exploration and management of us geological resources. *Engineering Science & Technology Journal*, 5(2), 313-337. <https://doi.org/10.51594/estj.v5i2.763>
- Pei, T., Xu, J., Liu, Y., Huang, X., Zhang, L., Dong, W., ... & Zhou, C. (2021). GIScience and remote sensing in natural resource and environmental research: Status quo and future perspectives. *Geography and Sustainability*, 2(3), 207-215. <https://doi.org/10.1016/j.geosus.2021.08.004>
- Ncube, M. M., & Ngulube, P. (2024). Enhancing environmental decision-making: a systematic review of data analytics applications in monitoring and management. *Discover Sustainability*, 5(1), 290. <https://doi.org/10.1007/s43621-024-00510-0>
- Terribile, F., Acutis, M., Agrillo, A., Anzalone, E., Azam-Ali, S., Bancheri, M., ... & Basile, A. (2024). The Land support geospatial decision support system (S-DSS) vision: Operational tools to implement sustainability policies in land planning and management. *Land Degradation & Development*, 35(2), 813-834. <https://doi.org/10.1002/ldr.4954>

- Hosen, B., Rahaman, M., Kumar, S., Sagar, L., & Akhtar, M. N. (2023). Leveraging artificial intelligence and big data for advanced spatial analytics and decision support systems in geography. *Malaysian Applied Geography*, 1(2), 62-67. <https://doi.org/10.26480/magg.02.2023.62.67>
- Dritsas, E., & Trigka, M. (2025). Remote sensing and geospatial analysis in the big data era: A survey. *Remote Sensing*, 17(3), 550. <https://doi.org/10.3390/rs17030550>
- Singh, A., Mehrotra, R., Rajput, V. D., Dmitriev, P., Singh, A. K., Kumar, P., ... & Singh, A. K. (2022). Geoinformatics, artificial intelligence, sensor technology, big data: emerging modern tools for sustainable agriculture. *Sustainable agriculture systems and technologies*, 295-313. <https://doi.org/10.1002/9781119808565.ch14>
- Liang, H., Zhang, Z., Hu, C., Gong, Y., & Cheng, D. (2023). A survey on spatio-temporal big data analytics ecosystem: resource management, processing platform, and applications. *IEEE Transactions on Big Data*, 10(2), 174-193. <https://doi.org/10.1109/tbdata.2023.3342619>