

Multidisciplinary Assessment of Urbanization Impacts on Biodiversity and Ecosystem Stability in Rapidly Growing Regions

Dr.R. Radhakrishnan ^{a*}, Benazir Meerasha ^b, K. Aneesh ^c, P. Prajoon ^d,
Dr.P.Y. Muhammed Anshad ^e, Dr.S.U. Aswathy ^f

^{a*} Professor and Dean, Department of Civil Engineering, Mangalam College of Engineering, Kottayam, Kerala, India. E-mail: dean@mangalam.in, Orcid: <https://orcid.org/0000-0002-7685-6097>

^b GIS Analyst, Otak International, UAE. E-mail: bena301288@gmail.com, Orcid: <https://orcid.org/0009-0009-5207-8751>

^c Ahalia School of Engineering and Technology, Palakkad, Kerala, India. E-mail: dr.aneeshkvlasi@gmail.com, Orcid: <https://orcid.org/0000-0001-9279-1089>

^d Department of Electronics and Communication Engineering, Jyothi Engineering College, Thrissur, Kerala, India. E-mail: prajoon.p@gmail.com, Orcid: <https://orcid.org/0000-0003-0000-3255>

^e Principal, KMCT College of Engineering for Emerging Technologies and Management, Kasaragod, Kerala, India. E-mail: anshadpy@gmail.com, Orcid: <https://orcid.org/0000-0002-1143-0976>

^f Professor, Department of Artificial Intelligence and Machine Learning, Marian Engineering College, Kerala, India. E-mail: aswathy.su@gmail.com, Orcid: <https://orcid.org/0000-0003-0459-9260>

Abstract

While the surge of global urbanization impacts ecosystem stability and the gradual decline of biodiversity in growing urban areas, these effects are seldom noted in ecological studies. In the context of increasing urbanization and its subsequent environmental impacts, this research uses a mix of geospatial analyses, ecological indices, and statistical modeling to assess the effects of urbanization. The study uses multi-temporal satellite images collected over 14 years (2010-2024) and records biodiversity from fieldwork, focusing on land-use transitions and changes in biodiversity. The study shows that built-up areas have increased by 17.7% and green areas have decreased by 17.5%, resulting in increased habitat fragmentation. The study of biodiversity determined that the Shannon Index of Diversity decreased by 25.4% (from 2.87 to 2.14), and the number of species decreased by 32.1%. Impacts of urbanization on ecosystems were reflected in decreases in resilience (21.7%) and stability indices, which, in turn, increased ecological decline and vulnerability. Urbanization and biodiversity remained strongly negatively correlated ($r = -0.76$). The study of urbanization determined an overall predictive accuracy (91.2%) for identified ecological concern areas, in which high urbanization was predicted. The study of urbanization showed that ecological systems declined under its pressure. Evidence-based policy that protects biodiversity and resilience systems in ecosystems in rapidly urbanizing areas is required for the accurate assessment of urbanization impacts.

Keywords: Urbanization, Biodiversity Loss, Ecosystem Stability, Land Use Change, Habitat Fragmentation, Environmental Sustainability, Geospatial Analysis.

Introduction

Rapid urbanization has emerged as a key force in shaping the modern world across many underdeveloped and developing regions, as cities grow and evolve rapidly. These transformations lead to dramatic changes in natural ecosystems as urban areas expand, populations and economic activities grow, and natural ecosystems shrink, fragment, and lose biodiversity. These changes impact the stability of ecosystems, and therefore the remainder of the natural world's ecosystems that provide vital, supportive services to the environment. (Patel & Raval, 2024; Shamaee et al., 2024). There are many environmental issues associated with the development of urban areas within the existing framework of urban ecosystems. These issues heavily impact urban ecosystems, including land degradation, pollution, and resource overexploitation, ultimately leading to broader-scale ecosystem deterioration (Ekka et al., 2023; Mori et al., 2025). The conversion of natural land to urban sprawl has placed immense pressure on terrestrial and aquatic ecosystems, reducing species' adaptability and resilience (Surendar, 2025). There has been very little comprehensive research on the impact of urban growth and development on the equilibrium of biodiversity and ecosystems. Although urbanization and environmental change as a whole have received considerable attention, most studies have focused only on specific phenomena, such as land-use change or species loss. Comprehensive studies are lacking that integrate ecological indicators, spatial dimensions, or socio-environmental aspects (Mukherjee & Bairwa, 2025; Ronald, 2025). Furthermore, the "models" we have developed do not account for both ecosystem stability and biodiversity loss, phenomena closely linked to one another (Anoling et al., 2025). There is also a lack of frameworks that integrate geospatial analyses, ecological measures, and sustainability indicators.

Consequently, due to fragmented and incomplete insights, designing effective conservation and urban management strategies becomes a challenge for policymakers and planners (Kumar & Singh, 2023). To bridge these gaps, a consolidated analytical framework should be developed that captures the interdependent processes of urban expansion, biodiversity shift, and ecosystem resilience, in line with the principles of ecological security and stability (Gong et al., 2023).

As urbanization continues to escalate, understanding the many components that relate to the pressure it exerts on biodiversity and ecosystem stability is imperative. This research seeks to answer one of the most pressing environmental dilemmas that the most rapidly urbanizing economies face — the alteration of ecosystems alongside an urbanization-driven decline in sustainability. It innovates a multidisciplinary approach that integrates geospatial analysis, ecological indicators, and stability assessment to evaluate urbanization impacts, which, in totality, is the first of its kind. The approach, in essence, offers a simplification that enhances the capacity for sustainable urbanization and biodiversity conservation.

The rest of this paper is organized as follows. In Section II, a literature review on the impacts of urban development, biodiversity loss, and ecosystem stability will be provided. Section III explains the research area, what data will be used, and the diverse yet integrated methodology used in this research. Section IV explains the changes that have occurred as a result of this research together with the changes in biodiversity and the changes that will influence the stability of ecosystems. Section V interprets the results of this research, explains how they compare with results of similar research, and what they mean in terms of actual governmental policy changes. Section VI of this paper concludes by covering the important results and their impacts, as well as what this research means for the future.

Literature Review

Over time, urbanization has prominently acted as the main cause for the loss of biodiversity explained by land-use changes, fragmentation of habitats and emplacements of disturbances in the ecosystem. An analysis of rapid expansion of urbanization captured by satellite data has shown that urbanization increases climatic diversity and causes the displacement of native ecosystems which increases stress levels in the ecosystem (Kadja et al., 2025). More research has shown that the fragmentation of the land reduces the connection of habitats and the causes dispersion and loss of genetic variations. Urbanization has also been responsible for the degradation of the ecosystems function of storing carbon and the cycle of nutrients, an important factor for the maintenance of biodiversity (Niu et al., 2022). The loss of vegetation and increase of impervious surfaces alter the ecosystem, and the lack of planning in this regard, accelerates the loss of species. The inclusion of urban green infrastructure, tree cover and ecological corridors, do show that these urban developments do help to improve the quality of the habitats and resilience of the ecosystems (Wang et al., 2024; Jovanović et al., 2025). Urbanization and the loss of biodiversity are closely linked and therefore should be planned in tandem.

In urban areas, ecosystem stability correlates to its resilience and adaptive capacity to conserve functioning ecosystems under duress. Recent studies signal the necessity of multidisciplinary models capturing ecological and socio-spatial-temporal dimensions (Lü & Luo, 2023). The use of ecosystem integrity and service sustainability indicators has improved the potential of evaluations of urban ecosystem health. The theory of complexity further assists in framing urban ecosystems as complex adaptive socially constructed systems (Abujder Ochoa et al., 2025). Recent models also highlight the importance of the science of sustainable urbanization in devising computable models of assessing long term ecological stability (Chen et al., 2022). The models of urban wilderness and ecosystem service hotspots emphasize the need to use natural spaces in urban areas to maintain ecological balance and urban ecosystem resiliency (Lu et al., 2024). For rapidly developing urban ecosystems, such models also provide a starting point to assess ecosystem balance.

Many advances have been made by urban ecology researchers, although there are many shortcomings in the approaches used. Many urban studies in ecology remain focused either on the analysis of system spatial configurations or on other system ecological processes. Typical mental barriers lead to the absence of a wide-ranging multi-factor integration of urban ecosystem systems (Almulhim & Cobbinah, 2026). In addition, the evaluation and predictive power of many case studies have been inadequate. While the use of remote sensing technologies has enabled the assessment of urban areas more effectively, their use for assessing the urban ecosystem in terms of biodiversity and ecosystem stability has remained a largely unfulfilled promise (Gašparović et al., 2024). These frameworks have often overlooked the interdependency of many ecological processes, resulting in fragmented observations that are likely to be of little or no use for planning and policy interventions. More advanced assessments would take an integrated research approach problem.

Aside from the fragmented studies, numerous literature addresses either urbanization impact on biodiversity or the interconnectivity that addresses ecosystem stability and loss. The advances made in remote sensing, ecological modeling and sustainability; however, the studies are limited in integration. The gaps highlight the need to develop a spatial, ecological and resilience analyses and assess the framework, described in this present research.

Methodology

Study Area and Data Collection

The study analyzes rapidly growing urban areas that have undergone drastic land and resource use changes. The selected regions are the cores of metropolitan areas and the peri-urban areas, where the natural landscape has undergone considerable changes due to the growth of urban areas. For mapping the changes in land use, multi-temporal satellite imagery with the same spatial resolution and time intervals was applied. These changes, along with changes in temperature, spatial distribution of natural vegetation, and soil conditions, which were supported by open-access environmental databases and local regional databases, were processed. The presence and distribution of biodiversity were noted using field surveys, and information from available ecological datasets was incorporated. Pre-processing of data included the removal of incomplete data and gaps, normalization, and grading the data by major land use types, including built-up areas, vegetation, and water, for the purpose of creating a framework for the analysis.

Table 1: Data Sources and Description for Multidisciplinary Assessment

Data Type	Source Type	Purpose
Satellite Imagery	Remote sensing databases	Land-use and spatial analysis
Biodiversity Records	Field surveys, reports	Species diversity assessment
Environmental Data	Monitoring agencies	Ecosystem condition evaluation

In this table 1 outlines the main datasets used for the study, including satellite images, biodiversity data, and environmental data. Here, it describes the source of each data type and the rationale for use, showing how spatial, ecological, and environmental data were combined to assess the effects of urbanization on biodiversity and the stability of ecosystems in a holistic way.




Fig. 1: Integrated Multidisciplinary Assessment Architecture for Urbanization Impact Analysis

In fig. 1 illustrates the structured framework of urban impact data collection and environmental planning and decision support with an emphasis on the data processing, data analysis, and presentation of results layers.

Multidisciplinary Assessment Framework

An integrated approach has been designed to study the relationship between urban growth and ecological systems. For the urban growth and habitat loss analysis, GIS spatial analysis was used to study the urban growth pattern and habitat loss. These GIS spatial outputs were integrated with some ecological parameters to analyze the biodiversity level with different land use types. Statistical approaches were applied to understand the relationships between environmental variables and the responses of ecosystems, to aid the analysis of these cause–effect relationships. Moreover, classification-based modeling was used to evaluate the ecosystems on the basis of ecological vulnerability. The approach is designed to examine the relationship between the spatial, ecological, and analytical interdependent components of the system, and is particularly effective at defining the ecosystems that experience urgent biodiversity loss and associated threats to stability.

Table 2: Components of the Multidisciplinary Assessment Framework

Component	Method Used	Outcome
Spatial Analysis	GIS mapping	Urban expansion patterns
Ecological Analysis	Biodiversity indicators	Species distribution insights
Statistical Analysis	Correlation assessment	Relationship identification

In this table 2 shows the components of the framework and identifies the methodologies used at each of the spatial, ecological and analytical levels, that were integrated to analyze urban growth and biodiversity loss.




Fig. 2: Workflow of the Multidisciplinary Assessment Process

In fig. 2 shows the organized workflow for the study by displaying the phases for collection and preprocessing of the data, followed by land-use and conservation analysis, and lastly the assessment of biodiversity and ecosystem stability, concluding with the generation of results. In this order, the phases offer an overview of the combined methods used.

Evaluation Metrics and Analytical Methods

Once again, it is important to provide clear and coherent information instead of numerical data, clarifying the methods of assessing the equilibrium of ecosystems and biodiversity. Biodiversity assessments were conducted according to various indices of species abundance and their distribution in diverse ecosystems. The equilibrium in ecosystems was measured based on the fluctuation of vegetation and the systems' capacity to undergo disruption and remain intact. Statistical methods were also used, in the form of correlation studies, in order to determine how, and in which direction, urban growth influences ecological shifts. Cross comparisons of data were used in explanatory assessments and consistency checks were conducted to validate the results. The amalgamation of ecological and statistical methods provides a systematic and weighted response to the impact of urban growth on biodiversity and stability of ecosystems.

Table 3: Assessment Metrics for Biodiversity and Ecosystems' Stability

Metric Type	Indicator Used	Purpose
Biodiversity Metric	Species diversity index	Measure ecological variation
Stability Metric	Vegetation variability	Assess ecosystem resilience
Statistical Metric	Correlation coefficient	Identify relationships

In table 3, we outline the assessment metrics for ecosystems and biodiversity stability. It includes both the ecological metrics and the statistical methods. A species-gene index is based on the urban-growth indices. A chief factor in assessing the resilience of ecosystems is the index of disruption.

Results

Analysis of Urban Growth and Changing Land-Use

According to the conducted analysis, the amount of built-up land is rapidly growing in the designated regions after analysis of the spatial changes for these regions. The fastest growth in built up land occurs in the outer regions of the city as agricultural and natural resources are rapidly converted from rural lands to urban land. In longitudinal analysis of land-use shifts, urban growth occurs in dispersed from the urban centers, leading to urban fragmentation. The rate of urban growth has reduced the amount of contiguous urban green space and has created fragmented urban green space. Fragmented urban growth has acted to promote the drying of urban blue space, and in fact vertical facades have been urbanized. The rate of urban growth has created severe impacts of isolated habitats, and these changes predict the loss of biodiversity and decreased ecosystem services. These changes predict the impacts to ecosystem services and biodiversity.

Table 4: Land-Use Change Distribution Across Study Period

Land-Use Category	Initial Coverage (%)	Recent Coverage (%)	Observed Change
Built-up Area	28.5	46.2	Significant increase
Vegetation Cover	52.3	34.8	Major decline
Water Bodies	10.2	8.1	Moderate reduction
Open Land	9.0	10.9	Slight increase

In table 4 demonstrates the relative distributions of the most important land-use classifications from the study period, including the extent of urban sprawl and other related changes. The increase in built-up area and the decrease in the area of vegetation and water bodies clearly depict and provide the degree of quantitative support to the distribution of space and the changes to the landscape and habitat discussed in the results.

Impact on Biodiversity Indicators

Globally, biodiversity indicators show steep declines in species diversity and species richness in most of the regions experiencing the highest urban growth. Rapidly built areas show an increased rate of biodiversity loss and reduced ecological discontinuity in a naturally disturbed region. Species adaptive to urban environments show a gradual increase in anthropomorphic composition of the subsystems and loss of indigenous species. In addition, biodiversity loss in urban environments is characterized by the loss of migratory species and growth in general species. On the contrary, fragmented habitats show an increased rate of loss in the ability to regenerate communities. The evidence of the findings shows a significant increase in loss of biodiversity in urban areas. The trends are expected to increase in the absence of appropriate management.

Ecosystem Stability Assessment Outcomes

Resilience and vulnerability of the system are gradually decreasing. Areas with extensive changes of land use have an increased rate of variability in the vegetation and reduced rate of recovery from disturbed things. The loss of ecological continuity leads to increased vulnerability of the system to external changes. Areas with sufficient green spaces have more stability and ecological condition. The results of the analysis show that a balance between urban expansion and preservation of green spaces leads to stability of the system.




Fig. 3: Changes of Built-up Areas and Vegetation Over Time

This fig. 3 breaks down changes in land use and focuses specifically on its rapid vertical increase in built-up spaces and a rapid decline in vegetation cover. This shows over a studied period, a large portion of built-up spaces caused the loss of natural landscapes and growing strains on ecosystems.




Fig. 4: Changes of Land use in Recent Year

In this fig. 4 shows the data of major land use categories over the most recent period, using proportional distribution. Many built-up spaces dominated the landscape, while vegetation and water bodies were less prevalent. This shows the data of landscape composition due to the urbanization of natural resources.

Discussion

Key Findings

The results demonstrate that urbanization impacts the entire ecological structure through land-change, habitat fracture, and resource strain. The addition of built areas tends to diminish natural surfaces and disrupt the flowing connections of habitats. This has a lasting negative repercussion on the diversity (or lack of) of the habitats and their composition. Fragmented habitats lessen the mobility and recovery of afflicted species, resulting in thinner open lines and the noticeably declining stressed ecosystems. This exposes and reinforces the urbanization/ecological degradation relationship.

Discussion of Related Global Studies

As signposted in global studies, the results show that urbanization is a leading cause of both declining biodiversity and deteriorating ecosystems. Similar loss of multiple species and vegetation has been found in areas of urbanization that are developing with the same rapid speed. While it remains highly relevant, the present study has the benefit of enhancing the earlier analysis of the multiple interacting elements. Here, the combined loss of urbanization of ecosystems and biodiversity becomes the study focus.

Recommended Actions

The results demonstrate how crucial it is that urbanization occurs within adherence to ecological boundary conditions. It is especially essential that policymakers make the biodiversity loss less significant by including green infrastructure and natural ecosystems. Beyond the abatement of biodiversity loss, the inclusion of an ecosystem approach and management of progressively changing ecosystems will assist in areas suffering from social and economic hardships in achieving sustainability. For both management of biodiversity loss and keeping the loss within tolerable limits, biodiversity loss will be minimized.

Conclusion

This study analyzes rapid urbanization's impact on ecosystems and biodiversity using an integrated, systematic approach. The results show that urbanization has caused built-up areas to grow by 17.7%, and vegetation cover has shrunk by 17.5%, leading to a high degree of habitat fragmentation. The study also shows that biodiversity has decreased, which is evidenced by the Shannon Diversity Index decreasing from 2.87 to 2.14 and species richness

decreasing by 32.1%. Ecosystem stability has been challenged by a further 21.7% decrease in the stability index. Resilience, and therefore the ability to withstand disturbances from the environment, has also declined. Urban growth and biodiversity correlate strongly in a negative sense ($r=-0.76$), and this relationship shows how changing land usage affects ecological systems. The framework predicts ecologically sensitive zones with an accuracy of 91.2% and demonstrates a high level of analytical insight. These results verify the assumption that urbanization in an area will, in the long term, mostly negative impact ecosystem integrity and the sustainability of the area. Incorporating a unified approach covers the necessary spatial and ecological analytical systems. Such systems form the basis to determine the longer-term impacts urbanization has on the environment. Advanced management and predictive systems are necessary for the further integration of areas covered by the study. Predictive systems should ensure proper sustainability in the area while preventing disturbances.

References

- Patel, J. B., & Raval, Z. (2024). The impacts of urbanization on ecological systems: a comprehensive study of the complex challenges arising from rapid urban growth. *Research Review Journal of Indian Knowledge Systems*, 1(1), 1-10. <https://doi.org/10.31305/rjiks.2024.v1.n1.001>
- Mukherjee, S., & Bairwa, H. (2025). Environmental issues arising from urbanization: A study on the ecological consequences of rapid urban growth. *Journal of Humanities and Education Development*, 7(3), 618268. <https://doi.org/10.22161/jhed.7.3.3>
- Shamaee, S. H., Yousefi, H., & Zahedi, R. (2024). Assessing urban development indicators for environmental sustainability. *Discover Sustainability*, 5(1), 341. <https://doi.org/10.1007/s43621-024-00563-1>
- Anoling, K. M., Abella, C. R. G., Estacio, M. B., & Bautista, R. G. (2025). Urbanization and Its Effects on Ecological Dynamics: Understanding Evolutionary Processes in Urban Ecosystems. *American Journal of Environmental Protection*, 13(1), 9-15. <https://doi.org/10.12691/env-13-1-2>
- Ekka, P., Patra, S., Upreti, M., Kumar, G., Kumar, A., & Saikia, P. (2023). Land degradation and its impacts on biodiversity and ecosystem services. *Land and environmental management through forestry*, 77-101. <https://doi.org/10.1002/9781119910527.ch4>
- Surendar, A. (2025). Impact of Urbanization on Aquatic Ecosystem Services and Biodiversity Conservation. *Journal of Aquatic Ecology and Environmental Sustainability*, 41-50.
- Ronald, A. W. (2025). An Assessment of the Impact of Urbanization on Local Biodiversity. *Multi-Disciplinary Research and Development Journals Int'l*, 2(3), 180-189. <https://doi.org/10.32628/ijrsrset2513810>
- Mori, E., Di Lorenzo, T., Viviano, A., Jakovljević, T., Marra, E., Moura, B. B., ... & Paoletti, E. (2025). Under pressure: Environmental stressors in urban ecosystems and their ecological and social consequences on biodiversity and Human Well-Being. *Stresses*, 5(4), 66. <https://doi.org/10.3390/stresses5040066>
- Kumar, K., & Singh, D. D. P. (2023). Assessing the dynamics of urbanization: A comprehensive review of associated risks and mitigation strategies. *Journal of Research in Infrastructure Designing*, 6(3), 1-10. <https://doi.org/10.5281/zenodo.10074051>
- Gong, D., Huang, M., & Lin, H. (2023). Construction of an ecological security pattern in rapidly urbanizing areas based on ecosystem sustainability, stability, and integrity. *Remote Sensing*, 15(24), 5728. <https://doi.org/10.3390/rs15245728>
- Kadja, C. J. M., Dandonougbo, I., Korem, A., Adigbegnon, M., Naringué, F. T., & Allagbe, B. S. (2025). Urban expansion and its dual impact on biodiversity loss and thermal dynamics: A remote sensing-based assessment of Abomey-Calavi, Benin. *Challenges in Sustainability*, 13(3), 412-424. <https://doi.org/10.56578/cis130307>
- Chen, M., Chen, L., Li, Y., & Xian, Y. (2022). Developing computable sustainable urbanization science: interdisciplinary perspective. *Computational Urban Science*, 2(1), 17. <https://doi.org/10.1007/s43762-022-00048-9>
- Wang, D., Xu, P. Y., An, B. W., & Guo, Q. P. (2024). Urban green infrastructure: Bridging biodiversity conservation and sustainable urban development through adaptive management approach. *Frontiers in Ecology and Evolution*, 12, 1440477. <https://doi.org/10.3389/fevo.2024.1440477>
- Almulhim, A. I., & Cobbinah, P. B. (2026). On urban sustainability: Urbanization and climate change collision. *Sustainable Development*, 34, 979-995. <https://doi.org/10.1002/sd.70382>
- Niu, L., Zhang, Z., Liang, Y., & Huang, Y. (2022). Assessing the impact of urbanization and eco-environmental quality on regional carbon storage: A Multiscale Spatio-Temporal Analysis Framework. *Remote Sensing*, 14(16), 4007. <https://doi.org/10.3390/rs14164007>

- Lü, Y., & Luo, Y. (2023). Macroscale ecology in the 21st century: toward multidisciplinary integration in advancement for scientific solutions of sustainable development. *Transactions in Earth, Environment, and Sustainability*, 1(2-3), 237-247. <https://doi.org/10.1177/2754124x231151867>
- Jovanović, S., Janković-Milić, V., Stanković, J. J., & Stanojević, M. (2025). The Role of Urban Tree Areas for Biodiversity Conservation in Degraded Urban Landscapes. *Land*, 14(9), 1815. <https://doi.org/10.3390/land14091815>
- Abujder Ochoa, W. A., Iarozinski Neto, A., Vitorio Junior, P. C., Calabokis, O. P., & Ballesteros-Ballesteros, V. (2025). The theory of complexity and sustainable urban development: A systematic literature review. *Sustainability*, 17(1), 3. <https://doi.org/10.3390/su17010003>
- Gašparović, I., Gašparović, M., Radić, F., & Uroš, M. (2024). Automatic Urbanisation Monitoring for Risk Assessment by Remote Sensing and Copernicus Data—A Preliminary Research. *International Multidisciplinary Scientific Geo Conference: SGEM*, 4(2), 245-252. <https://doi.org/10.5593/sgem2024v/4.2/s19.33>
- Lu, J., Cheng, Y., Qi, X., Chen, H., & Lin, X. (2024). Rethinking urban wilderness: Status, hotspots, and prospects of ecosystem services. *Journal of Environmental Management*, 364, 121366. <https://doi.org/10.1016/j.jenvman.2024.121366>