International Academic Institute for Science and Technology

International Academic Journal of Social Sciences
Vol. 1, No. 2, 2014, pp. 87-90.
ISSN 2454-3918

International Academic Journal of Social Sciences

Addiction and psychological factors related to the ethology of social crime prevention

Seyed Abolqasem Seyedan, Reza khodashenas abdolabade, najme khodashenas abdolabade

professor of psychology, Torbat Heydarieh universities, Iran.
Student -shahed rajaei torbat heydareye, Farhangian University, Tehran, Iran.
Psychologist, Teacher, Iran.

Abstract

Addiction is one of the major social harms and is the cause for the highest number of imprisonments in Iran. Thus the study of the causes influencing its creation plays a major role in the control, prevention and cure of other social phenomena.

Considering the fact that the role of family and social institutions like training institutions cannot always follow the individual plus hereditary (genetic) and environmental coefficients impacting addiction are generally out of the control of social workers and therapist, it is necessary that the psychological factors – as the most controllable factors – affecting addiction be taken into account. And the share of the individual's will power and mindset, such as identity formation, excitement, mood and will power be studied.

Thus the objective of the present research has been to identify the said factors of addiction. It is hoped that our therapists will lay emphasis on these factors and find the most controllable factor in this field.

Keywords: Addiction; Psychological factors addiction, Personality, excitation

Introduction:

Addiction as the main factors associated with crime and other social ills known and its role in the formation of abnormal social phenomena such as theft, divorce, drug trafficking, extortion and clearly seen and proven. Since the control factor involved in addiction like Family structure, social environment and the fabric of people's lives, proximity to drug-producing countries and … The control therapists and social workers and psychologists outside the treatment area. So be aware of the role of the individual in the field and Effective psychological analysis of its features and upgrades psychological ability to control and treat the phenomenon of the action.

The main body of the text or article

Addiction is a great danger that threatens teenagers and young adults. In our country the control of the family and the school is more intense in adolescence with fewer young people are facing addiction. Unfortunately, the failures of the last stage of adolescence in school and university entrance exams Unemployment and uncertainty about the future makes a group of teenagers with family background and

undermining belief to be driven by addiction. Age of addiction and its causes in our country is different from Western countries and that's why the solution of this problem as well different solutions classical school of Western psychology is about people addicted to their communities. It is often thought that a small amount of drugs, For example, smoking is safe. Of course, this works for a group of teenagers a curious action and manifestation of adults know themselves but not always the case. Use the minimum number of cigarettes for teens that there are other problems A prelude to the more vile behaviour and in some cases, Drug addiction is progressive and can be stronger in the future for drug use Such as alcohol, cannabis, heroin, opium and even the ground. Especially in the first and second stages of adolescence, the period of distance and separation of supply (Ages 12-18 years) the risk for all teens (Lotfabade).

Risk of addiction among teenagers and young adults are not the same. Adolescents and young people that the mood among the people "demand sensitive" (Ross Barrett, 1989) are threatening risk of addiction. Treat these people with intense emotions and because drug use is their emotional needs likely to be drawn into it. In addition, patients suffering from attention deficit disorder and conduct disorder hyper stimulation the risk of addiction threatens.

The factors leading to addiction can be divided into five categories:

1. Pharmaceutical and pharmacological agents 2. Psychological Factors 3. Family factors
4. Social factors 5. Political factors

Of course, we can say that any of the factors mentioned in connection with other factors. But in this article we'll discuss in particular Psychological factors.

Psychological factors, people addicted in one of the following five characters:

A) Normal person that female drug due to the indiscriminate use of analgesic medication was.

B) To eliminate their anxiety drug neurotic refuge.

C) Saykopat people to create ecstatic and happy, taking addictive substances.

D) Saykopat people who take drugs for depression.

E) Sykopaty combined with the characteristics and neurotic people (Ahmade).

Theoretical Foundations

A) According to psychoanalytic perspective in all human beings, there is a principle called the pleasure principle. The natural tendency of the organism to avoid pain and Search through the release of tension is fun. Freud's pleasure principle in connection with the finds Id. Id instinct is irrational and immoral and function. The first stage of psychosexual development child that is eighteen months in the name of oral stage most had a role in the formation of addiction in adulthood. At this point all the fun your child's oral gratification. Oral cause stress scenarios Children in search of satisfaction of mouth Characterized by calm after care and stroked. Exclusion or persuasion too may, Cause pathological traits in adults. Many traits can be mentioned that this step comes, that play a role in shaping addiction. The people at this stage in adults due to consolidation, following oral gratification are the most adults have things they must seek oral gratification. Drugs due to meet the security features that are capable of satisfying the individual's enjoyment. It is also possible for those who are strict superego and If your are punished to fix the guilt and stress, unconsciously turn to alcohol to get rid of the pressure of the superego.

B) Biological perspective: As we know, based on the orientation of the face, Addiction is merely due to exposure to opiates. When a person experiences stress, endorphins are released and A lifter stress caused numbness and pain tolerance increases. Some researchers have suggested that the use of heroin and other opiates may the collapse of long-term biochemical machine that makes endorphins. According to this view, because the drug addict natural pain reliever systems in cluttered his body continues to consume opiates (Sarason and Sarason). In addition to the above, the biological point of view from another dimension can be linked to activity in the nucleus accumbens by asking pointed out. Addicted to actually increase the amount of what you want not necessarily how you like it. In other words, when people become addicted to something, it is the dominant focus and stronger nucleus accumbens responds to it. Frequent consumption of cocaine release of dopamine in the nucleus accumbens its ability to and as well

as its ability to activate the part of the right prefrontal cortex and Search the drug's tendency to increase (Robinson and Bridge, 2001).

C) The behaviorist views, such addiction can be explained by conditioning factors: Any action that rewards practice it is learned to be followed. In the process of drug addiction makes people feel happy and joyful. The consent by the part of the brain called the reward pathway in the brain occur and this increase is treated. Like other addictive behavior is learned in this approach and spends the learning process. In general, in view of radical behavior such as Skinner, Everyone amplifiers to control the behavior controlled. In view of methodological behaviorists such as Julian Rotter and Bandura, Everyone has role models in the community control, controlled behavior (Laura Berk, 2007).

Characteristics and thrill-seeking addicts

2 initial period of 7 years of life, Individual character builds and with this interpretation have highlighted the role of families in this area, he said. Experience has shown that a drug addict to get his illegal and immoral acts and Freed from the shackles of moral crimes as well. Neurotic, psychotic personalities in the whole of Venice can be very vulnerable to addiction, he said (Lotfabade).

The role of emotional intelligence in addicts

People with substance abuse with no real understanding of their emotions the ability to correct behavior, logical and based on the reality of the day, On the other hand, these people also have difficulty in applying the right emotions. In general, explain the psychological features of the new plan Angle Emotional Intelligence That can have a large number of applications. Because unlike intelligence, emotional intelligence is not fixed and immutable trait, Can be used in rehabilitation centers and drug treatment center to rehabilitate these people's emotional intelligence.

Addiction social contribution in the incidence of injury

Divorce: One of the known causes of divorce is tendency to addiction, because, in these circumstances, children lose your guide to the immaturity, easily influenced by other people are (Laura Berk, 2007).

Domestic violence: To reverse the do's and do not for a woman or a husband, wife abuse is said to be and in all classes, from the poorest to the richest and educated families happen that addiction is one of the most well-known causes of domestic violence.

Of the most important ways to prevent and reduce domestic violence can be referred to drug treatment. Thus, to minimize or prevent lowering agents and prophylactic drug abuse should be identified.

Theft: White and colleagues (1985) found that alcohol, marijuana and hard drugs and petty crime and crime related property. The relationship between substance abuse and committing various criminal acts show that there is a link between certain combinations of materials and mass. Crack, break your consumers prefer to earn money, because immediately concludes. Although heroin users prefer night time theft, because their need is less urgent. (Bennett and Holloway, 2005).

Prostitution: The cause of many social pathologies and addiction disorders, family and individual. Especially the increasing phenomenon of addiction and dependence and addiction new considering the role emotional, mental, moral and social, family and community threatened and the pathogenic behavior, especially in cases of drug addiction and prostitution leads women to sexual deviation.

Addicted women compared with other women, have more psychological problems. It is known that depression or anxiety disorders the most common disease among women addicted to drugs. Studies also show that women with a history of sexual abuse were addicted, Most have experienced deep depression and semen taste As well as a high level of anxiety has been diagnosed, While addicted men with antisocial personality disorder marked (Hojjati).

Suicide: After mental disorders, substance abuse, suicide is the second largest risk factor and in both permanent and severe associated with suicide. This is related to the effects of Stupor and promiscuity many psychoactive drugs when someone like mourning and grief associated with grief, It increases the risk of suicide. Also, most people who commit suicide when they under the influence of sedative-hypnotic drugs (such as alcohol, benzodiazepines) are and In general, countries with higher levels of

alcohol consumption, higher suicide rates that the issue is primarily related to the alcohol used for alcohol distillation.

Cocaine and methamphetamine abuse is highly correlated with suicide. Among those who use cocaine, especially in the most at-risk youth in "consumption" or "left" on the, While in this role is older alcoholism.

Some mental disorders associated with alcohol abuse and the suicide rate is very high in alcoholic. Alcoholic depression and those who abuse alcohol, is very high. Long-term use or abuse of prescription benzodiazepines, as well as permanent as that associated with depression is associated with suicide (Savin- Williams and Ream 2003).

Conclusion:

What happened is part of social problems, crime and drug addiction was their direct effects. Considering the proven contribution in the formation of drug crimes large and small and shaping it to the top of the prison, we can say: Way out of many social problems such as divorce, suicide, sale and distribution of drugs and the prevention and treatment of addiction.

However, due to poor handling of environmental issues, social, biological and genetic therapists and social workers and specialists in the behavioural sciences, Factors and psychological factors can contribute the most effective way with other prevention methods And combating addiction and the damage caused by it. So, be aware of the role of the individual in this field and analyses the psychological characteristics affecting it and psychological knowledge to control and treat this phenomenon was acting.

References:

Ahmadi, A. Psychology of Adolescents and Youth (sixteenth edition). Isfahan: Minor publications

Arms den, G.C, McCauley. , Greenberg,M.T., Burke,P,& Mitchell.(1991). Parent and peer attachment in early adolescence depression.

Bennett. And Holloway. (2007), Drug-Crime Connections, University of Glam organ: Pontypridd.

Burke, Laura (2007). Developmental Psychology (from adolescence to the end of life) / (Volume II). Translation: Sayed Mohammad Yahiya. Tehran: Publication Arasbaran.

Gizzi, M.C. and Gerkin, P.(2010), Methamphetamine Use and Criminal Behavior, International Journal of Offender Therapy and Comparative Criminology, 54, 915-936.

Harrison.L.D.(1992), The Drug-Crime Nexus in the USA, Contemporary Drug Problems, 19, 181-202.

Islam, like Soraya (1392). Addiction (etiology and treatment) / (fifth edition). Tehran: PNU.

Lotfabade, H. Developmental Psychology (2): Youth and adult (twelfth edition). Tehran: Publication side.

Rio, John Marshall (2001). motivation and excitement. Translation: Yahya Syed Mohammad (1382). Tehran: Publication editing.

Wertheim U, Degkwitz P, Karausz M. Mental disorders and the course of opiate dependence. Psychiatry Prax 2000; 27(2): 77-85.